

Observations of groundreflected Global Navigation Satellite System signals from space

Clara Chew

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA © 2017. All rights reserved.

Outline 5

- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work

Outline J


- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work

Monostatic Radar


Bistatic Radar


Measures backscatter


Measures forward scatter












Monostatic Radar


Measures backscatter


Bistatic Radar


Measures forward scatter











Monostatic Radar


Bistatic Radar


Measures backscatter


Measures forward scatter
















- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work

Sources of spaceborne reflections

1. Soil Moisture Active Passive (SMAP) radar receiver:


- H and V ~36 dB
- Data since Aug 2015
- Polar orbiting


2. TechDemoSat-1 (TDS-1):

- LHCP antenna ~13 dB
- Data every 8 days since July 2014
- Polar orbiting


3. Cyclone GNSS (CyGNSS):

- LHCP antenna ~13 dB
- Launched Dec 2016, no data yet
- Orbits the tropics











What do data look like recorded over the <u>ocean</u>?




- Very rough: Incoherent scattering
- Large spatial footprint: 25 km

What do data look like recorded over land?



- If smooth, coherent scattering
- Variable footprint: 0.5 25 km




Outline J

- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work







- Mosaic Cropland/Natural Ve
- Forest or Shrubland
- Regularly Flooded Soil
- Urban
- Water




= Regions of High SNR (dB)








Punjab region of India and Pakistan

Agricultural region
Heavily irrigated during growing season
Crops sown in November, harvested in April












Do this just for soil moisture




Outline - 5


- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work









Summer: Jun, Jul, Aug 2016




Winter: Dec, Jan, Feb 2015/2016







Changes of collocated reflections over time



Changes of collocated reflections over time



Modeled changes due to F/T



Outline 5

- Forward vs. backscatter
- Spaceborne GNSS-Reflectometry data
- Sensitivity to near-surface soil moisture
- Sensitivity to landscape freeze/thaw state
- Future work



jpl.nasa.gov